

CONN
S
43
.E22
NO.957



PLEASE HANDLE
WITH CARE

University of
Connecticut Libraries



3 9153 01231318 7

*The
Connecticut
Agricultural
Experiment
Station,
New Haven*

New Tobacco Cyst
Nematode Resistant
Shade Tobacco
Varieties

BY JAMES A. LAMONDIA

Bulletin 957

May 1999

Two varieties of Connecticut shade tobacco (*Nicotiana tabacum* L.) resistant to the tobacco cyst nematode, *Globodera tabacum tabacum* (Lownsbery & Lownsbery) Stone, have been developed at The Connecticut Agricultural Experiment Station's Valley Laboratory. These varieties, 'Poquonock' and 'Metacommet', allow shade tobacco production in cyst nematode-infested soils without the annual use of fumigant nematicides. Yields and sorting qualities are equal to or better than the nematode-susceptible shade variety 'O-40'. Limited quantities of seed are available to growers and scientists.

The tobacco cyst nematode was first discovered in Hazardville, CT in the early 1950's (Lownsbery and Lownsbery, 1954). It is now widely distributed in shade production areas in the Connecticut River Valley of Connecticut and Massachusetts. A related nematode, *Globodera tabacum solanacearum* (Miller and Gray, 1972) Stone, suppresses the growth and yield of tobacco in Virginia and North Carolina (Komm et al., 1983; Melton et al., 1991). Nematode infection of roots can cause dramatic stunting, yield loss, and reduce leaf quality. A nonlinear damage function was developed to predict shade tobacco yield losses based on preplant nematode densities in soil (LaMondia, 1995). Losses can exceed 40 percent in Connecticut at high nematode densities. Nematode losses have been prevented by the use of preplant fumigation of soils. However, fumigant nematicide use is expensive and the number of fumigants available for use has been reduced due to environmental considerations.

BREEDING AND SELECTION

All Connecticut shade and broadleaf varieties tested were susceptible to *G. t. tabacum* (LaMondia, 1991). Flue-cured tobacco lines with resistance to *G. t. solanacearum* were found resistant to *G. t. tabacum* (LaMondia, 1988), and crosses to Connecticut shade tobacco were begun in 1987. Resistance to *G. t. tabacum* was later determined to be conferred by a single dominant gene (LaMondia, 1991).

Both Poquonock and Metacommet are inbreds derived from a cross between the nematode-susceptible Connecticut shade tobacco line 'O-30' and the nematode-resistant flue-cured line VA-81. Resistance to *G. t. tabacum* and *G. t. solanacearum* may be linked to wildfire (*Pseudomonas tabaci* Wolf & Foster, Stevens) resistance (Spasoff, 1971). Wildfire resistance was transferred from *Nicotiana longiflora* to the breeding line TL 106, which had a pair of chromosomes from the wild species (Clayton, 1947). *Nicotiana longiflora* was resistant to *G. t. solanacearum* in pot experiments (Baalawy and Fox, 1971). Metacommet was selected and inbred over 12 selfed generations of field evaluation for horticultural characteristics and greenhouse evaluation for *G. t. tabacum* resistance. Poquonock was

selected after backcrossing to Connecticut types to incorporate the resistance gene into a predominantly Connecticut shade background. The original O-30 and VA-81 cross was back-crossed to O-30 twice, back-crossed to the nematode-susceptible shade line O-40 twice, then crossed again to a selfed inbred (selfed for three generations) of O-30 and VA-81. The resulting line was selfed to homogeneity (stable inbred genetics) over six generations. For both varieties, individual plants in the F2 and F4 generations were selected for cyst nematode resistance using a greenhouse seedling assay (LaMondia, 1991). Progeny testing was performed in 1993 to select plants homozygous for *G. t. tabacum* resistance.

Poquonock and Metacommet both contain the dominant hypersensitive gene for resistance to tobacco mosaic virus derived from *Nicotiana glutinosa* L. Both lines have also been selected for reduced sensitivity to weather fleck caused by ozone. Flue-cured tobacco varieties with resistance to *G. t. solanacearum* were intolerant of infection, and required nematicide use to maintain yields of resistant plants (Komm et al, 1983). We selected Poquonock and Metacommet under field conditions for growth and yield in the presence of damaging population levels of *G. t. tabacum* to avoid severe problems with intolerance to nematode infection.

EVALUATION AND PERFORMANCE

The effects of nematode resistance on *G. t. tabacum* populations were determined in field plots in a cloth-covered shade tent at the Experiment Station's Valley Laboratory in Windsor in 1993, 1994, and in 1997. Plants were grown in four rows in replicate 25m² plots. Nematode densities were determined from 25 composite core samples to 15 cm deep taken prior to planting and again after harvest. Cysts were recovered from soil by flotation in a modified Fenwick can, crushed, and the number of juveniles in eggs counted. Changes in density were expressed as the ratio of final to initial counts (Pf/Pi). Nematode resistant varieties reduced population densities in soil by approximately 70% (Table 1).

Table 1. The effects of nematode resistant or susceptible shade tobacco varieties on *G. t. tabacum* populations in naturally infested soils.

Variety	Reaction	Pf/Pi ratio*		
		1993	1994	1997
O-40	Susceptible	2.91	2.97	3.01
Metacommet	Resistant	0.17	0.26	0.28
Poquonock	Resistant	ND	0.33	0.37
P=		0.001	0.001	0.001

* Nematode population change expressed as the ratio of final to initial densities.

Table 2. Yield and quality characteristics of *G. t. tabacum* resistant or susceptible tobacco varieties.

Variety	Reaction	Yield/plant (g)		Quality Grade (%)1996*			\$ Value per ⁺ 45 kg 1993
		1993	1996	1	2	3	
O-40	Susceptible	466.1	614.7	46	20	33	1,088
Metacommet	Resistant	746.3	759.6	26	36	38	1,902
Poquonock	Resistant		731.9	44	43	13	1,980
P=		0.09	0.001				

* Grade 1 = excellent wrapper quality, grade 3 = poor.

⁺ Economic value determined by price per grade and cured leaf weight, 1993.

Over the same time period, nematode populations increased by about 200% in soil where a susceptible variety grew. The population reduction resulting from the season-long production of a resistant variety is comparable to growing a nematode-susceptible variety with soil fumigation (LaMondia, 1993).

Leaf yield and quality of Metacommet, Poquonock and the nematode-susceptible O-40 were compared in field plots infested with 120 to 250 infective *G. t. tabacum* juveniles per cm³ soil (Table 2). Fresh weight yields were similar or greater for nematode-resistant varieties compared to the O-40 standard. Cured leaf quality was determined from samples submitted to commercial shade tobacco producers for curing and evaluation. Quality of Metacommet and Poquonock, as determined by percent weight in each grade in 1996, and by economic value of yields in 1993, was comparable to or better than O-40.

Metacommet and Poquonock shade varieties will allow the production of high quality shade tobacco in fields infested with damaging populations of *G. t. tabacum*. These varieties give growers a non-chemical control which will reduce nematode populations as well as a fumigant nematicide.

REFERENCES

- Baalawy, H.A., and J.A. Fox. 1971. Resistance to Osborne's cyst nematode in selected *Nicotiana* species. *Journal of Nematology* 3:395-398.
- Clayton, E.E. 1947. A wildfire resistant tobacco. *Journal of Heredity* 38:35-40.
- Komm, D.A., J.J. Reilly, and A.P. Elliot. 1983. Epidemiology of a tobacco cyst nematode (*Globodera solanacearum*) in Virginia. *Plant Disease* 67:1249-1251.
- LaMondia, J.A. 1995. Shade tobacco yield loss and *Globodera tabacum tabacum* population changes in relation to initial nematode density. *Journal of Nematology* 27:114-119.
- LaMondia, J.A. 1993. Evaluation of reduced fumigation rates on tobacco cyst nematode populations and shade tobacco yield. *Fungicide and Nematicide Tests* 48:216.
- LaMondia, J.A. 1991. Genetics of tobacco resistance to *Globodera tabacum tabacum*. *Plant Disease* 75:453-454.
- LaMondia, J.A. 1988. Tobacco resistance to *Globodera tabacum*. *Annals of Applied Nematology* 2:77-80.
- Lownsbery B.F. and J.W. Lownsbery. 1954. *Heterodera tabacum* n. sp., a parasite of solanaceous plants in Connecticut. *Proceedings of the Helminthological Society of Washington* 21:42-47.
- Melton, T.A., J.A. Phillips, J.L. Imbriani, and K.R. Barker. 1991. First report of *Globodera tabacum solanacearum* on flue-cured tobacco outside Virginia. *Plant Disease* 75:1074.
- Miller, L.I., and B.J. Gray. 1972. *Heterodera solanacearum* n. sp., a parasite of solanaceous plants. *Nematologica* 18:404-413.
- Spasoff, L., J.A. Fox, and L.I. Miller. Multigenic inheritance of resistance to Osborne's cyst nematode. *Journal of Nematology* 3:329-330.

ACKNOWLEDGMENTS

The author wishes to thank Dr. C.S. Johnson for seed of *G. t. solanacearum*-resistant lines, H.F. Brown, Inc. and Culbro, Inc. for excellent cooperation in cured leaf quality analysis, and J. Canepa-Morrison and R. Horvath for technical assistance.

The Connecticut Agricultural Experiment Station is an Equal Opportunity, Affirmative Action Employer. Persons with disabilities who require alternate means of communication of program information should contact the Station Editor at (203) 974-8446 (voice); (203) 974-8502 (FAX); or paul.gough@po.state.ct.us (E-mail)



The Connecticut Agricultural Experiment Station, founded in 1875, is the first experiment station in America. It is chartered by the General Assembly to make scientific inquiries and experiments regarding plants and their pests, insects, soil and water, and to perform analyses for State agencies. The laboratories of the Station are in New Haven and Windsor; its Lockwood Farm is in Hamden. Single copies of bulletins are available free upon request to Publications; Box 1106; New Haven, Connecticut 06504.

ISSN 0097-0905



University of
Connecticut
Libraries



39153029109099

